

Dactyliopius adonidum.  
 " bromiliæ.  
 " citri.  
 " destructor.  
 Diaspis amygdali.  
 " pyricola.  
 " rosæ.  
 Eriococcus araucariæ.  
 " coriaceous.  
 " eucalypti.  
 Eulecanium persicæ.  
 Fiorinia camelliae.  
 Icerya Purchasi.  
 Lecanium cerasarum.  
 " depressum.  
 " hemisphaericum.  
 " hesperidum.  
 " longulum.  
 " oleæ.  
 Mytilaspis citricola.  
 Parlatoria zizyphus.  
 Pseudolecanium tokionis.  
 Pulvinaria camellicola.  
 Anthonomus pomorum (apple-blossom weevil).  
 Aphis mali (apple-aphis).  
 " persicæ (green peach-aphis).  
 " prunifoliae (plum-aphis).  
 Aspidictus perniciosus (*San José scale*).  
 Blastodacna hillerae and vinolentella (apple-pith moth).  
 Byturus tomentosus (raspberry-beetle).  
 Chrysobothris femorata and C. mali (apple-tree borer).  
 Diplosis pyriversa (pear-midge).  
 Doryphora decemlineata (*Colorado beetle*).  
 Euthrips fuscus (*thrips*).  
 " pyri (*thrips*).  
 Hedyia ocellana (apple-bud moth).  
 Hoplocampa testudinea (apple saw-fly).  
 Hylemyia antiqua (onion-fly).  
 Incurvaria capitella (currant-shoot borer).  
 Lampronia rubiella (raspberry-moth).  
 Leptops Hopei (apple-root borer).  
 Mytilaspis pomorum (mussel or oyster scale).  
 Myzus cerasi (black aphid).  
 Nysius vinitior (*Rutherford bug*).  
 Otiorrhynchus picipes (raspberry-weevil).  
 Phorodon humuli (hop-aphid).  
 Psylla mali (apple-sucker).  
 Pulvinaria vitis (woolly currant-scale).  
 Rhopalosiphum ribis (currant-aphid).  
 Rhizopertha collaris (apple-tree borer).  
 Saperda candida (round-headed tree-borer).  
 Schizoneura lanigera (*American blight*).  
 Scolytus rugulosus (fruit-bark beetle).  
 Sesia tipuliformis (currant-clearwing).  
 Siphonophora rosæ (rose-aphid).

—  
THIRTEENTH SCHEDULE.  
(DESTRUCTION.)

Anarsia lineatella (peach-moth).  
 Bacillus amylovorus (pear-blight or fire-blight of the pear).  
 Citrus canker of Japan.  
 Daucus cucumeris.  
 " facialis.  
 " Kirki.  
 " melanotus.  
 " passifloræ.  
 " psidii (*New Caledonian fruit-fly*).  
 " raratiogensis.  
 " tongæ.  
 " virgatus.  
 " xanthodes (pine-apple fruit-fly).  
 Halterophora capitata (*Mediterranean or West Australian fruit-fly*).  
 Heliothis obsoleta (tomato-caterpillar).  
 Laverna herellara.  
 Lonchæa splendida (tomato fruit-fly).  
 Merodon equestris (narcissus-fly).  
 Rhyzoglyphus echinopus (bulb-mite).  
 Tephritis Tryoni (*Queensland fruit-fly*).  
 Trypetis musæ (*New Hebrides fruit-fly*).  
 " pomonella (apple-maggot).

—  
FOURTEENTH SCHEDULE.  
(DIPPING.)

Actinonema rosæ (rose black-blotch).  
 Armillaria mellea (stringy-rot).  
 Ascochyta Boltshauseri (broad-bean anthracnose).  
 " pisi (pea-spot).

Bryobia pratensis (red mite).  
 Cladosporium fulvum (tomato-leaf blight).  
 Clasterosporium carpophilum (peach die-back).  
 Colletotrichum Lindemuthianum (French-bean anthracnose).  
 Cylindrosporium chrysanthemi (chrysanthemum-leaf blight).  
 Eriophyes pyri (pear-mite).  
 " ribis (currant gall-mite or big bud).  
 Exoascus deformans (peach-curl).  
 Fusarium limonis (collar-rot (citrus)).  
 Macrosporium nobile (carnation-rust).  
 " tomato (tomato black-spot).  
 Microspheara grossulariae (gooseberry-leaf mildew).  
 Nematus ribesi (gooseberry saw-fly).  
 Ocneria dispar (gypsy-moth).  
 Oosporia scabies (potato-scab).  
 Peronospora cubensis (cucumber and melon mildew).  
 " Schleideni (onion-mildew).  
 Phoma omnivora (wither-tip, foot-rot of citrus trees).  
 " sanguinolenta (carrot-disease).  
 Phragmidium rubi-idei (raspberry-rust).  
 " subcorticatum (rose-rust).  
 Phylosticta apii (celery-leaf spot).  
 " prunicola and circumissa (shot-hole fungus).  
 Podosphaera leucotricha (apple-mildew).  
 Puccinia asparagi (asparagus-rust).  
 " bullata (celery-rust).  
 " hieracii (chrysanthemum-rust).  
 " pruni (prune and peach rust).  
 Septoria ribis (currant-leaf spot).  
 " rosæ (rose-leaf spot).  
 Sphaerotheca castagnæi (hop-mildew).  
 " mors-uvæ (gooseberry-mildew).  
 " pannosa (rose-mildew).  
 Sporodesmium griseum (grey scab of lemon).  
 Uromyces appendiculatus (haricot-bean rust).  
 " betæ (beet-rust).  
 " fabæ (broad-bean rust).  
 Vermicularia circinans (onion-scab).  
 " varians (black-dot of potato, tomato, &c.).

—  
FIFTEENTH SCHEDULE.

(RESHIPMENT OR DESTRUCTION.)

Aegeria tipuliformis (currant or gooseberry borer).  
 Alternaria solani (early blight).  
 Bacillus phytophthorus (potato black-leg).  
 " solanacearum (potato-bacteriosis).  
 " sp. (potato wet-rot).  
 Bacterium hyacinthi (bulb-bacteriosis).  
 " tumefaciens (hairly-root, crown-gall, or root-knot).  
 Bitter pit.  
 Cacococcia postvittana (light-brown apple-moth).  
 Carpocapsa pomonella (codlin-moth).  
 Coniothecium chomatosporum.  
 Dendrophagus globosus (crown-gall).  
 Eumerus strigatus (lesser narcissus-fly).  
 Fusarium citri (verrucosis of lemon).  
 " lycopersici (sleepy disease of tomato).  
 " oxysporum (potato dry-rot).  
 Fusicladium dendriticum (apple-scab).  
 " pyrinum (pear-scab).  
 Gloeosporium venetum (raspberry-spot).  
 Glomerella rufo-maculans (bitter-rot).  
 Heterosporium echinulatum (carnation fairy-ring).  
 Hypholoma fasciculare (raspberry root-knot).  
 Lemon bacteriosis.  
 Lita solanella (potato-moth).  
 Melanoze.  
 Monilia fructigena (ripe-rot).  
 Nectria cinnabarina (canker).  
 " ditissima (canker).  
 Phoma citricarpa (anthracnose or black-spot of orange).  
 Phyloxxera vastatrix (vine-louse or phylloxera).  
 Phytophthora infestans (potato-rot or Irish potato-blight).  
 Plowrightia morbosæ (black-knot).  
 Polystictus cinnabarinus (bark-rot).  
 Pseudomonas campestris (cabbage black-rot).  
 Pythiacystis citrophthora (lemon brown-rot).  
 Rhizoctonia solani (potato collar-fungus).  
 Samenoidea opalescens (peach-root borer).  
 Sclerotinia bulborum.  
 " sclerotiorum (sclerotium).  
 Sphaerella brassicæcola (cabbage-leaf spot).  
 " tabifica (beetroot and mangold rot).  
 Synchytrium endobioticum (wart-disease of potatoes).  
 Tylenchus devastatrix (eelworm).

J. F. ANDREWS,  
 Clerk of the Executive Council.