

- Avena fatua* L. (Black or Wild oat).
Brassica spp.
Bromus mollis L. (Soft Brome grass).
Bromus secalinus L. (Chess or Cheat).
Camelina sativa Crantz (False flax).
Carex spp.
Caucalis spp.
Cenchrus spp.
Chenopodium album L. (Fat Hen).
Chondrilla juncea L. (Chondrilla).
Cichorium Intybus L. (Chicory).
Convolvulus spp.
Crepis spp.
Crotalaria sagittalis L. (Rattle Box).
Cryptostemma calendulaceum R.Br. (Cape-weed).
Cyperus rotundus L. (Nut grass).
Datura spp.
Diodia teres Walt. (Button-weed).
Dipsacus sylvestris L. (Wild Teazel).
Echium vulgare L. (Blue weed).
Eichornia speciosa Kunth. (Water Hyacinth).
Emex australis Steinh. (Spiny Emex).
Eragrostis major Host. (Stink grass).
Erechtites valerianifolia D.C. (Brazilian Fire-weed).
Eruca sativa Mill (Rocket).
Eryngium spp.
Erysimum repandum L. (Treacle Mustard).
Euphorbia spp.
Foeniculum vulgare Mill (Fennel).
Galium aparine L. (Cleavers or Goose grass).
Hedypnois cretica Wild (Hawkbit or Cretica).
Helminthum autumnale L. (Sneeze weed).
Homeria collina Vent. var. *minuta* Sweet (Cape Tulip).
Hordeum murinum L. (Barley grass).
Iva axillaris Pursh. (Poverty weed).
Iva xanthifolia Nutt. (Marsh Elder, False Sunflower).
Jatropha spp.
Juncus spp.
Lantana spp.
Lactuca spp.
Lappula spp.
Lithospermum arvense L. (Corn Gromwell, Iron-weed).
Leontodon hirtus L. (Hairy Thrinacia).
Linum spp.
Lolium temulentum L. (Drake or Darnel).
Lychnis Flos-cuculi L. (Ragged Robin).
Lychnis Githago Scop. (Purple Corn Cockle).
Madia sativa Mol. (Pitch weed).
Marrubium vulgare L. (Horehound).
Matricaria spp.
Melilotus officinalis Lam. (Yellow Melilot).
Nicotiana glauca Grahm. (Wild Tobacco plant or Tree Tobacco).
Panicum sanguinale L. (Summer grass).
Panicum colonum L. } Barnyard Grasses.
Panicum Crus-galli L. }
Papaver spp.
Picris spp.
Polygonum spp.
Ranunculus spp.
Raphanus Raphanistrum L. (Wild Radish or Jointed Charlock).
Rapistrum spp.
Reseda luteola L. (Wild Mignonette).
Rhus radicans L. (Poison Ivy, Poison Oak, or Poison Vine).
Romulea spp.
Rumex spp.
Salvia sylvestris L. (Wild Sage).
Salvia verbenaca L. (Purple Sage).
Saponaria spp.
Scirpus spp.
Senecio spp.
Setaria glauca Beauv. (A Pigeon grass).
Setaria verticillata Beauv. (A Pigeon grass).
Setaria viridis Beauv. (A Pigeon grass).
Sherardia arvensis L. (Field Madder).
Sida spp.
Silene spp.
Sisymbrium spp.
Solanum spp.
Spergula spp.
Stachys arvensis L. (Stagger-weed).
Tagetes spp.
Thlaspi arvense L. (Wild Garlic, Bastard Cress, French Weed).
Tribulus spp.
Urtica spp. ~~Urtica~~
Verbascum spp.
Verbenae spp.
Xanthium spp.

SCHEDULE D.

Agricultural Seeds.	Weed Seeds allowed in addition to those previously enumerated.	Percentage (maximum) of such Weed Seeds allowed.
<i>Lolium perenne</i> (Perennial rye-grass)	Italian rye-grass (<i>Lolium italicum</i>)	10
<i>Trifolium repens</i> (White clover)	<i>Trifolium dubium</i>	5
Timothy	Red clover .. White clover ..	5 3
Cocksfoot	Perennial rye-grass Italian rye-grass	3 3
Italian rye-grass ..	Perennial rye-grass	15

SCHEDULE E.

Agricultural Seeds.	Maximum Percentage (by Weight) of Impurities permitted.	Minimum Percentage of Germination.
Asparagus	$\frac{1}{2}$	55
Barley	$\frac{1}{2}$	80
Beans (broad)	$\frac{1}{2}$	70
Beans (French)	$\frac{1}{2}$	75
Beans (Mauritius)	$\frac{1}{2}$	60
Beans (Soy)	$\frac{1}{2}$	60
Beans (others)	$\frac{1}{2}$	60
Beets	$\frac{1}{2}$	60 (of clusters)
Borecole	$\frac{1}{2}$	60
Brocoli	$\frac{1}{2}$	60
Brussels Sprouts	$\frac{1}{2}$	60
Buckwheat	$\frac{1}{2}$	75
Cabbage	$\frac{1}{2}$	60
Cauliflower	$\frac{1}{2}$	60
Carrot	$\frac{1}{2}$	55
Celery	$\frac{1}{2}$	50
Chicory	$\frac{1}{2}$	60
Clovers, medics, &c.—		
<i>Lespedeza striata</i> (Japanese clover)	$\frac{1}{2}$	40
<i>Lotus angustissimus</i> (Lotus—Narrow-leaved)	$\frac{1}{2}$	60
<i>Lotus corniculatus</i> (Birdsfoot trefoil)	$\frac{1}{2}$	60
<i>Lotus hispidus</i> (Boyd's clover)..	$\frac{1}{2}$	60
<i>Lotus major</i> (Greater Birdsfoot trefoil)	$\frac{1}{2}$	60
<i>Medicago denticulata</i> (Burr clover)	$\frac{1}{2}$	70
<i>Medicago lupulina</i> (English trefoil)	$\frac{1}{2}$	70
<i>Medicago orbicularis</i> (Button clover)	$\frac{1}{2}$	50
<i>Medicago scutellata</i> (Snail clover)	$\frac{1}{2}$	60
<i>Melilotus alba</i> (Bokhara or Sweet clover)	$\frac{1}{2}$	35
<i>Trifolium alexandrinum</i> (Egyptian clover)	$\frac{1}{2}$	65
<i>Trifolium fragiferum</i> (Strawberry clover)	$\frac{1}{2}$	40
<i>Trifolium hybridum</i> (Alisks clover)	$\frac{1}{2}$	75
<i>Trifolium incarnatum</i> (Crimson clover)	$\frac{1}{2}$	75
<i>Trifolium pratense</i> (Red clover)	$\frac{1}{2}$	75
<i>Trifolium pratense</i> var. <i>perenne</i> (Perennial red, Chilian, and Cow-grass clover)	$\frac{1}{2}$	75
<i>Trifolium repens</i> (white clover)	$\frac{1}{2}$	70
<i>Trifolium subterraneum</i> (Subterranean clover)	$\frac{1}{2}$	70
Cotton	1	70
Cowpeas	$\frac{1}{2}$	75
Cress, Garden (<i>Lepidium sativum</i>)	$\frac{1}{2}$	60
Cress, Water (<i>Nasturtium officinale</i>)	$\frac{1}{2}$	50
Cucumber	$\frac{1}{2}$	70
Egg Plant (<i>Solanum Melongena</i>)	$\frac{1}{2}$	50
Endive	$\frac{1}{2}$	55
Grasses—		
<i>Agrostis</i> spp. (Red-top, Creeping Bent, Rhode Island Bent, Fiorin)	$\frac{1}{2}$	50
<i>Alopecurus pratensis</i> (Meadow Foxtail)	$\frac{1}{2}$	50